



HOW TO USE YOUR PENILE INJECTABLE MEDICATION



FOLLOW THESE DIRECTIONS, UNLESS OTHERWISE DIRECTED BY YOUR PHYSICIAN

Your compounded prescription drug will be injected into the spongy tissue of the penis. The spongy tissue runs the length of the penis on the right and left sides



Choose an injection site on one side of the shaft of the penis. **Avoid any veins you can see.** With each use change the side of the penis you inject. Also change the spot you inject.



You should sit up straight or lie back slightly when you inject the drug. Hold the head of your penis with your thumb and first finger. Then stretch your penis lengthwise along your thigh and you can clearly see the spot you want to inject. Clean the injection spot with a new alcohol swab. Do not throw this swab away. You will need to use it again.



Hold the penis firmly against your thigh to keep it from moving during the injection. Hold the syringe between your thumb and first finger. Then push the needle into the injection site as far as it will go.



Push the plunger rod as far as it will go to inject the whole dose. If the solution does not flow easily, move the needle a little bit and push as before. Pull the needle out of your penis. Hold the alcohol swab on the injection site for about 5 minutes or until any bleeding stops. Carefully put the outer cap back on the needle. Dispose of used supplies safely. Your pharmacist may be able to supply a box made for needles.

The information provided is not intended to cover all possible uses. This information is generalized and not intended as specific medical advice. If you have questions about the use of your prescription, please contact your prescriber.