

NSAID Containing Transdermal Gels

NSAIDS Containing PLO Gels: NSAIDS stands for non-steroidal anti-inflammatory drugs. These gels contain the active ingredients found in brand products such as: Advil™, Nuprin™, Aleve™, Orudis™, Actron™, Motrin™, Ketoprofen, Ibuprofen. These gels may also be combined with locally acting anesthetics or products for neuropathic pain.

Common Uses: This medication is usually used to treat inflammation and to provide analgesia or eliminate pain.

How To Use This Medication: This product may be supplied either in a tube (with or without measuring device), or in calibrated syringes, and should be applied to clean skin.

Precautions: Do not exceed the recommended dosage set by your doctor. This product contains lecithin. Patients should not use this medication if they are allergic or hypersensitive to NSAIDS. Persons allergic or sensitive to egg whites may have a reaction to this product. You should avoid the use of aspirin and other NSAIDS (Advil™, Nuprin™, Aleve™, Orudis™, Actron™, Motrin™, Ketoprofen, Ibuprofen) while on this medication. This medication should be stored out of the reach of children.

Possible Side Effects: The product may cause local irritation at the site of administration after it is applied. If this occurs, patients should discontinue use of the medication and notify their physician or pharmacist. Patients may also experience a feeling of warmth or coolness. This is considered normal and does not indicate an allergic reaction.

If You Miss a Dose: If you are taking this medication on a regular basis, and miss a scheduled dose, apply it as soon as you remember, however do not double up doses.

The information in this monograph is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. The information is generalized and is not intended as specific medical advice. If you have questions about the drugs you are taking, check with your prescriber or pharmacist.