

## TETRACAINE MEDICATED LOLLIPOPS

**TETRACAINE LOLLIPOPS:** This specially made medication contains tetracaine in a sugar free base.

**Common Uses:** This medication is a local anesthetic and is used to numb a sore or irritated mouth or throat. It may also be used to numb the gag reflex in certain types of dental procedures and to relieve pain associated with oral lesions.

**How To Use This Medication:** Suck on the lollipop for 10 to 15 seconds, then stop. The medication should begin to work in approximately 2 minutes. This lollipop should be dissolved in the mouth and the saliva that is produced should be swallowed. Swallowing the saliva will numb the throat, keeping the saliva in your mouth for as long as possible will numb your mouth and tongue. The lollipop may also be moved around in the mouth for maximum results. This may be repeated until results are obtained and may be repeated every 2 to 3 hours or as directed by the your prescriber.

**Precautions:** Do not exceed the recommended dosage set by your doctor. **KEEP THIS MEDICATION OUT OF THE REACH OF CHILDREN.** Do not break the pop and swallow the pieces of medication. *If used in children an adult should monitor the child at all times.* Follow the instructions in the section above “How to use this medication”. *The child and some adults may not need the entire lollipop at one time.*

**Possible Side Effects:** This medication may cause tingling, stinging, tenderness. If they persist, contact your doctor. Also, if you have any of the following, let your doctor know dizziness, lightheadedness, skin rash, tremors, or muscle aches.

This medication was specially prepared for you at Bedford Pharmacy.

If you have any questions, please feel free to call us.